



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

438. *By Prof. F. H. Loud.*—Two equiangular parallelograms, $OACB$ and $OA'C'B'$ are so placed that the equal angles AOB and $A'OB'$ coincide. The sides of the former figure are constant, those of the latter are variable, subject to the condition $A'O + OB' : AO + OB :: \text{area } OA'C'B' : \text{area } OACB$. $A'C'$ meets BC in D , and $B'C'$ meets AC in E . Show that DE passes through a fixed point, and determine the point.

439. *By Prof. Nicholson.*—Solve geometrically the following:

On a line whose length is a are two points x distance apart; what is the average value of x ?

440 *Selected.*—A lamina is bounded on two sides by two similar ellipses, the ratio of the axes in each being m , and on the other two sides by two similar hyperbolas, the ratio of the axes in each being n . These four curves have their principal diameters along the coordinate axes. Prove that the product of inertia about the coordinate axes is

$$\frac{(a^2 - a'^2)(\beta^2 - \beta'^2)}{4(m^2 + n^2)}$$

where aa' , $\beta\beta'$ are the semi-major axes of the curves. (Routh's Rigid Dynamics.)

PUBLICATIONS RECEIVED.

Celestial Charts, made at the Litchfield Observatory of Hamilton College, Clinton, N. Y., by C. H. F. PETERS. Nos. 1-20. Each chart covering 20 min. in right ascension and 5° in declination.

The whole of the work involved in the production of this very valuable series of charts, including the cost of publication, has been at the author's expense for gratuitous distribution.

Studies in Logic, by Members of the Johns Hopkins University. 203 pp. 12mo. Boston: Little, Brown, and Company. 1883.

Scientific Proceedings of the Ohio Mechanics' Institute. No. 1 of Vol. II., March, 1883. Cincinnati, Ohio.

An Extension of the Theorem of the Virial, and its Application to the Kinetic Theory of Gases. By H. T. EDDY C. E., PH. D. [From the Scientific Proceedings of the Ohio Mechanics' Institute for March, 1883.]

Report on the Michigan Forest Fires of 1881. [Signal Service Notes, No. I.] Office of Chief Signal Officer. Washington. 1882.

ERRATA.

- On page 74, line 4, for "1076.4", read ± 1076.4 . [“positive”.]
 “ “ 75, lines 12 and 13 from bottom, *dele* parenthesis before “in” and after
 “ “ 85, line 12, for “2”, read 1.